



**New Synthetic Methodology For The  
Ethyl[2-(2*H*-Chromene-3yl)-4-Oxo-1,3-Thiazolidin-3-yl]Acetates**

**N. Lakshmana Rao<sup>1</sup>, S. Kiran Kumar<sup>2</sup>, D. Rambabu<sup>3</sup>, M. V. Basaveswara Rao<sup>4\*</sup>**

1. Department of Chemistry, CMJ University, Shillong -793303, Meghalaya, **INDIA**
2. Department of Chemistry, K. L. University, Guntur-522502, Andhra Pradesh, **INDIA**
3. Department of Chemistry and the National Institute for Biotechnology in the Negev,  
Ben-Gurion University of the Negev, Be'er-Sheva 84105, **ISRAEL**
4. Department of Chemistry, Krishna University, Machilipatnam-521001, A.P, **INDIA**

Email: [vbrmandava@yahoo.com](mailto:vbrmandava@yahoo.com)

Received on 21<sup>st</sup> November and finalized on 22<sup>nd</sup> November 2013

---

**ABSTRACT**

*2H-3-chromene carbaldehydes react with glycine ethyl ester hydrochloride salt, and mercapto acetic acid in the presence of diisopropylethylamine to give ethyl [2-(2H-chromene-3yl)-4-oxo-1,3-thiazolidin-3yl]acetates. The products obtained in good yields and mechanism of this reaction was discussed.*

**Keywords:** 2*H*-3-chromene carbaldehydes, Thiazolidinones, DIPEA, mercapto acetic acid and 2-(2*H*-chromene-3yl)-4-oxo-1,3-thiazolidin-3yl]acetates.

---