



Facile Synthesis, Characterization of Novel Schiff Bases and N-Nucleosides Bearing Quinazoline Moiety and Evaluation of Their Antimicrobial Effects

M. A. EL-Hashash¹, M. A. Kadhim^{2*} and S. A. Rizk¹

1. Department of Organic Chemistry, Faculty of Science, University of Ain Shams, Cairo, **EGYPT**

2. Department of Chemistry, Faculty of Educational Science, University of Anbar, **IRAQ**

Email: mak.organic@yahoo.com

Accepted on 26th October 2015

ABSTRACT

The present work describes convenient synthesis of the novel Schiff bases **4**, **5** and **6** by reaction of each of the previously synthesized quinazolinones **1**, **2** and **3** with *p*-methoxybenzaldehyde. [2+2] Cycloaddition Reaction of the former Schiff bases within phenylisothiocyanate affording azatedine derivatives **7** and **8**. Also, the quinazolines **1** and **2** could be reacted with α -bromoglucose tetra acetate giving peracetylated N-glycosides of quinazolinones **9** and **10** which were then deacetylated to afford N-glycosylated quinazolinones **11** and **12** respectively. On the other hand, Dapsone was reacted with *p*-methoxyacetophenone to afford 4-[(4-aminophenyl) sulfonyl]-N-[(1E)-1-(4-methoxyphenyl)ethylidene]aniline **15** which was, in turn, reacted with 4-chloro-2-ethoxyquinazoline **16** affording Schiff base **17**. The latter was reacted with 2-furoyl chloride affording Schiff base **18**. The structures of the novel Schiff bases and N-glycosides were confirmed by the IR, ¹H-NMR, ¹³C-NMR, MS and elemental analysis. The antimicrobial activity for these synthesized compounds could outline, the moderate activity was observed with new quinazoline compounds which proved to possess marked activity against *E. coli*, *S. aureus* and *C. albicans*. The strong activity has observed nearly with the most synthesized compounds.

Keywords: quinazolin-4(3H)-one, N-glycosylated quinazoline, Dapsone, Schiff base, Azatedine and Arylidine.

INTRODUCTION

Quinazoline as N-heterocycles have received considerable attention in the literature as a consequence of their exciting biological properties and their role as pharmacophore[1-4]. Some of quinazolinones were screened in vitro for their antimicrobial activity and the energy gap between HOMO and LUMO has been calculated to reflect the chemical reactivity and kinetic stability of compounds [5]. A Novel series of N-substituted-2-phenylquinazolin-4-ones bearing different anilines at the N-3 of quinazolin-4-one scaffold via acetyl-flexible linker as anticancer agents with the compounds were synthesized by insertion of methylene (CH₂) bridge at C4-position of quinazoline moiety to provide a flexibility that increase their anti-proliferative activity against three human tumor cell[6]. Similarly, heterocycles containing the quinazoline moiety are of interest because they show some pharmacological and biological activities [7-9]. Quinazoline derivatives were reported to possess anticonvulsant[10], antitumor[11,12], antihyper

tensive[13], antithrombotic[14], antidiabetic[15], antitrypanosomal[16], anti-inflammatory[17]. Olaparib (Anticancer Agent), Zopolrestat and Ponalrestat (Anti-diabetic Agent), Azelastine and Flezelastine (Anti-histaminic Agent) could have benzyl moiety directly attached to quinazoline precursors. From this view the authors could be decided and prompted us to synthesize the quinazoline derivatives bearing benzyl moiety to evaluate their antimicrobial effects.

MATERIALS AND METHODS

All melting points recorded are uncorrected. The IR spectra were recorded on a Pye Unicam SP 1200 spectrophotometer using the KBr wafer technique. The ^1H - and ^{13}C -NMR spectra were determined on a Varian FT-200, Bruker AC-200 MHz spectrophotometry experiment using TMS as an internal standard. Chemical shifts (δ) are expressed in ppm. The mass spectra were determined by MP model NS-5988 and Shimadzu single focusing mass spectrometer (70 eV). All solvents used were of HPLC/AnalaR grade.

General Procedure for the Synthesis of the quinazoline-4(3H)-one 1-3: A mixture of 3,1-Benzoxazin-4-one (0.01 mol) and *p*-phenylene diamine, 4,4'-benzidine and/or 2,2'-dimethyl,4,4'-benzidine (0.01 mol) was heated under reflux in absolute ethanol (20 mL) for 6h. The resulting mixture was cooled and then poured onto an ice/water mixture. The separated solid was filtered, washed with water, air-dried and crystallized from ethanol to give colorless crystals of products.

2-(4-Aminophenyl)-4-(4-methoxy)benzylquinazolin-4(3H)-one (1): m.p. 168-170 °C; yield 76 %; Anal. for $\text{C}_{22}\text{H}_{19}\text{N}_3\text{O}_2$ (m.w. 357). Anal: Calcd: C, 73.94; H, 5.32; N, 11.76; Found: C, 73.93; H, 5.34; N, 11.75. IR ν (cm^{-1}) 1666 (C=O), 1610 (C=N), 2993 (CH); MS: m/z (int. %) $[\text{M}+\text{H}]^+$ 357(27.9); ^1H -NMR (DMSO- d_6) δ 3.22 (s, 3H, OCH_3), 4.02 (s, 2H, CH_2Bz), 6.3(s, 2H, NH_2), 7.18-7.92 (m, 12 H, 3Ph-H); ^{13}C -NMR: 15.01, 55.79, 114.41, 114.39, 120.81, 122.48, 122.51, 126.59, 126.68, 127.31, 128.72, 129.59, 129.62, 130.18, 131.18, 133.41, 146.92, 147.58, 154.01, 160.02, 160.60, 162.90.

2-(4-Amino-[1,1'-biphenyl]-4-yl) 4-(4-methoxy)benzylquinazolin-4(3H)-one (2): m.p. 182-184 °C; yield 68 %; Anal. for $\text{C}_{28}\text{H}_{23}\text{N}_3\text{O}_2$ (m.w. 433), Calcd: C, 77.59; H, 5.31; N, 9.69; Found: C, 77.57; H, 5.34; N, 9.69. IR ν (cm^{-1}) 1668 (C=O), 1622 (C=N), 2992 (CH). MS: m/z (int. %) $[\text{M}^+]$ 433(35.7); ^1H -NMR (DMSO- d_6) δ 3.41 (s, 3H, OCH_3), 4.04 (s, 2H, CH_2Bz), 6.23(s, 2H, NH_2), 6.95-7.90 (m, 16H, PhH); ^{13}C -NMR: 15.01, 64.91, 114.41, 120.82, 122.01, 122.69, 125.11, 126.19, 126.61, 126.72, 127.32, 128.69, 130.21, 130.21, 131.01, 131.01, 132.12, 133.39, 134.42, 134.78, 137.21, 140.11, 146.89, 150.01, 154.00, 160.58, 160.00, 162.91.

2-(4-Amino-3,3'-dimethyl -[1,1'-biphenyl]-4-yl)-4-(4-methoxy)benzylquinazolin-4(3H)-one (3): m.p. 188-190 °C; yield 68 %; Anal. for $\text{C}_{30}\text{H}_{27}\text{N}_3\text{O}_2$ (m.w. 461), Calcd: C, 78.09; H, 5.85; N, 9.11; Found: C, 78.07; H, 5.84; N, 9.09. IR ν (cm^{-1}) 1668 (C=O), 1622 (C=N), 2992 (CH). MS: m/z (int. %) $[\text{M}^+]$ 461(6.6); ^1H -NMR (DMSO- d_6) δ 2.19, 2.41 (2s, 2H, 2CH_3 of biphenyl), 3.84 (s, 3H, $-\text{OCH}_3$), 4.52 (s, 2H, $-\text{CH}_2\text{Bz}$), 6.95-7.50 (m, 4H, PhH), 7.33-8.10 (m, 6H, biphenyl), 7.48-8.19 (m, 4H, qui); ^{13}C -NMR: 15.01, 17.59, 18.61, 55.80, 114.41, 120.82, 122.01, 122.69, 125.11, 126.19, 126.61, 126.72, 127.32, 128.69, 130.21, 130.21, 131.01, 131.01, 132.12, 133.39, 134.42, 134.78, 137.21, 140.11, 146.89, 150.01, 154.00, 160.58, 160.00, 162.91.

General Procedure for the Synthesis of the Schiff Bases 4-6: A mixture of compounds 1, 2, and/or 3(0.01 mol) and *p*-methoxybenzaldehyde (0.01 mol) was heated under reflux in absolute ethanol (20 mL) for 6 h. The resulting mixture was cooled and then poured onto an ice/water mixture. The separated solid was filtered, washed with water, air-dried and crystallized from ethanol to give colorless crystals of products.

4-(4-Methoxy)benzyl -2-(4-((4-methoxybenzylidene) amino) phenyl)quinazolin-4(3H)-one (4): m.p. 200-202 °C; yield 80 %; Anal. for C₃₀H₂₅N₃O₃ (m.w. 475). Anal: Calcd: C, 75.78; H, 5.26; N, 8.84; Found: C, 75.76; H, 5.24; N, 8.85. IR ν (cm⁻¹) 1666 (C=O), 1610 (C=N), 2993 (CH); MS: m/z (int. %) [M⁺] 475 (40); ¹H-NMR (DMSO-d₆) δ 3.64, 3.82 (2s, 6H, 2OCH₃), 4.26 (s, 2 H, CH₂Bz), 7.18-7.78 (m, 12 H, 3Ph-H), 7.34-8.19 (m, 4 H, qui), 8.68 (s, 1H, =CH-); ¹³C-NMR: 15.01, 55.79, 64.89, 114.41, 114.49, 120.81, 122.48, 122.51, 126.59, 126.68, 127.31, 134.39, 140.81, 142.48, 142.51, 146.59, 148.68, 149.31, 151.72, 153.59, 155.62, 160.18, 161.18, 163.41, 164.92, 167.58, 168.01, 170.02, 170.60, 172.90.

4-(4-Methoxy)benzyl-2-(4'-(4-methoxybenzylidene)amino)-[1,1'-biphenyl]-4-yl)quinazolin-4(3H) -one (5): m.p. 212-214 °C; yield 74 %; Anal. for C₃₆H₂₉N₃O₃ (m.w. 551), Calcd: C, 78.40; H, 5.26; N, 7.62; Found: C, 78.37; H, 5.24; N, 7.59. IR ν (cm⁻¹) 1668 (C=O), 1622 (C=N), 2992 (CH). MS: m/z (int. %) [M⁺] 551 (35.7); ¹H-NMR (DMSO-d₆) δ 3.62, 3.84 (2s, 6H, 2OCH₃), 4.52 (s, 2H, CH₂Bz), 6.95-7.50 (m, 8H, PhH), 7.33-8.10 (m, 8H, biphenyl), 7.48-8.19 (m, 4H, qui), 8.72 (s, 1H, =CH-); ¹³C-NMR: 15.01, 17.59, 18.61, 55.80, 64.91, 114.41, 114.41, 120.82, 122.01, 122.69, 125.11, 126.19, 126.61, 126.72, 127.32, 128.69, 130.21, 130.21, 131.01, 131.01, 132.12, 133.39, 134.42, 134.78, 137.21, 140.11, 146.89, 150.01, 154.00, 160.58, 160.00, 162.91, 167.58, 168.01, 170.02, 170.60.

4-(4-Methoxy)benzyl-2-(4'-(4-methoxybenzylidene) amino)-3,3'-dimethyl -[1,1'-biphenyl]-4-yl) phth -alazin-1(2H)-one (6): m.p. 222-224 °C; yield 70 %; Anal. for C₃₈H₃₃N₃O₃ (m.w. 579), Calcd: C, 78.75; H, 5.69; N, 7.52; Found: C, 78.73; H, 5.67; N, 7.50. IR ν (cm⁻¹) 1668 (C=O), 1622 (C=N), 2992 (CH). MS: m/z (int. %) [M⁺] 579 (35.7); ¹H-NMR (DMSO-d₆) δ 2.19, 2.41 (2s, 6H, 2CH₃ of biphenyl), 3.64, 3.84 (2s, 6H, 2OCH₃), 4.52 (s, 2H, CH₂Bz), 6.95-7.50 (m, 8H, PhH), 7.33-8.10 (m, 6H, biphenyl), 7.48-8.19 (m, 4H, phthala), 8.72 (s, 1H, =CH-).

Synthesis of Compounds 7-8: Equimolar amounts of **4** and/or **5** (0.01 mol) and phenyl isothiocyanate in 25 mL toluene was refluxed for 6 h. The solvent was distilled off and the residue was washed with ethanol followed by water, and the product was crystallized from ethanol as yellow crystals.

2-(4-[2-(4-Methoxyphenyl)-3-phenyl-4-thioxo-1,3-diazetidino-1-yl]phenyl)amino-2-(4-methoxy benzyl)quinazolin-4(3H)-one (7): m.p. 278-280 °C, yield 68 %. Anal. for C₃₇H₃₀N₄O₃S (m.w. 610); Calcd: C, 72.78 ; H, 4.91; N, 9.18; S, 5.24; Found: C, 72.76; H, 4.87; N, 9.15; S, 5.23; IR ν (cm⁻¹) 1319 (C=S), 1519 (C=N), 1669 (C=O), 3318, 3443 (CH); MS: m/z (int. %) [M⁺] 610 (59.1); ¹H-NMR (DMSO-d₆) δ 3.46, 3.76 (2s, 6H, 2OCH₃), 4.37 (s, 2H; CH₂Bz), 6.70-7.15(m, 8H, 2C₆H₄OCH₃), 7.18-7.30(m, 5H, -C₆H₅), 7.32-7.40(m, 4H, X = Ar), 7.44-8.20 (m, 4H, qui).

2-(4-[2-(4-Methoxyphenyl)-3-phenyl-4-thioxo-1,3-diazetidino-1-yl]-[1,1'-biphenyl])amino-2-(4-methoxybenzyl)quinazolin-4(3H)-one (8): m.p. 284-286 °C, yield 72 %. Anal. for C₄₃H₃₃N₄O₃S (m.w. 685); Calcd: C, 75.32; H, 4.81; N, 8.17; S, 4.67; Found: C, 75.30; H, 4.83; N, 8.18; S, 4.69; IR ν (cm⁻¹) 1322 (C=S), 1522 (C=N), 1671 (C=O), 3327, 3433 (CH); MS: m/z (int. %) [M⁺] 685 (66.3); ¹H-NMR(DMSO-d₆) δ 3.52(s, 3H, OCH₃), 3.76 (s, 3H, -OCH₃), 4.45 (s, 2H; CH₂Bz), 7.17-7.33 (m, 5H, -C₆H₅), 6.70-6.95 (m, 8H, 2C₆H₄OCH₃), 7.0, 7.28(m, 8H, X =Ar), 7.30-8.20 (m, 4H, qui).

Synthesis of Acetylated Derivatives 9 and 10 and Deacetylated Derivatives 11 and 12: A crude mixture of quinazolinones **1** and/or **3** (0.05 mol) and α -bromoglucose tetraacetate (0.01 mol) in 100 mL 1, 4-dioxane was heated with frequent stirring under reflux for 4h. The mixture was cooled to room temperature and the solvent was removed under reduced pressure. The residue was dissolved in ethyl acetate, washed sequentially with saturated NaHCO₃ and dried over MgSO₄. The purification and separation was achieved by column chromatography (3:1 EtOAc: Hexane) giving the acetylated products **9** and **10** as a white solid which was later on recrystallized using dichloromethane- diethyl ether – hexane solvents. A solution of **9** or **10** in MeOH (50 mL) was treated with sodium carbonate solution (0.05 mol). A white solid syrupy began to settle. The residue was then purified by column chromatography (3:1,

EtOAc:Hexane) giving the deacetylated derivatives each as a white solid, which was crystallized by dichloromethane-diethyl ether-hexane solvents affording **11** or **12** respectively.

(2R, 3R, 4S, 5R, 6S)-2-(Acetoxymethyl)-6-((4-(4-methoxybenzyl-1-oxoquinazolin-2-yl) phenyl) amino) tetrahydro-2H-pyran-3, 4, 5-triyl triacetate (9): m.p. 124- 126 °C; yield 58 %. Anal. for C₃₆H₃₇N₃O₁₁ (m.w. 687): Calcd: C, 62.88; H, 5.38; N, 6.11; Found: C, 62.84; H, 5.38; N, 6.08. IR ν (cm⁻¹) 1662, 1738 (C=O), 2988 (CH), 3334 (NH); MS: m/z (int. %) [M⁺] 687(1.4); ¹H-NMR (DMSO-d₆) δ 2.05-2.06 (s, 12H, 4Ac-H), 3.45(s, 3H, OCH₃), 4.00 (s, 1H, sec NH), 4.52 (s, 2H, CH₂Bz), 3.75-5.11 (m, 6H, H-2, H-3, H-4, H-5, H-6a, H-6b), 5.22 (d, 1H, H_{anom}), 6.70-6.95 (m, 4H, C₆H₄OCH₃)7.01-7.34(m, 4H, X=Ar), 7.47-8.19 (m, 4H, qui); ¹³C-NMR: 15.01, 20.70, 21.02, 21.02, 21.02, 62.39, 64.89, 69.01, 69.38, 73.0, 77.18, 87.89, 117.89, 117.89, 120.8, 121.11, 122.42, 122.42, 126.59, 126.71, 127.32, 130.38, 130.8, 131.11, 132.42, 132.42, 136.59, 136.71, 143.19, 146.90, 154.01, 160.58, 170.20, 170.20, 170.21, 170.21.

(2R,3R,4S,5R,6S)-2-(Acetoxymethyl)-6-((4'-(4-methoxybenzyl-1-oxoquinazolin-2-yl)-3,3'-dimethyl-[1, 1'-biphenyl]-4-yl)amino)tetrahydro-2H-pyran-3,4,5-triyl triacetate (10): m.p. 164-166 °C; yield 56%. Anal. for C₄₄H₄₅N₃O₁₁ (m.w. 789): Calcd: C, 66.92; H, 5.70; N, 5.32; Found: C, 66.93; H, 5.70; N, 5.30. IR ν (cm⁻¹) 1666, 1736 (C=O), 2992 (CH), 3334 (NH); MS: m/z (int. %) [M⁺] 789(10.9); ¹H-NMR (DMSO-d₆) δ 2.05-2.12 (s, 12H, 4 Ac-H), 2.29, 2.22 (2 s, 6H, 2CH₃), 3.22 (s, 3H, -OCH₃), 4.0 (s, 1H, sec-NH), 4.52 (s, 2H, CH₂Bz), 3.74-5.14 (m, 6H, H-2, H-3, H-4, H-5, H-6a, H-6b), 5.47 (d, 1H, anom-H), 6.66-7.72 (m, 10H, ArH), 7.50-8.19 (m, 4H, qui); ¹³C-NMR: 15.01, 17.61, 17.89, 20.72, 21.01, 21.01, 21.01, 62.39, 64.90, 69.01, 69.39, 73.01, 77.18, 88.21, 113.89, 120.78, 122.01, 125.11, 125.68, 126.59, 126.72, 126.72, 127.29, 130.00, 131.00, 131.62, 133.41, 134.38, 134.78, 137.20, 140.78, 142.01, 145.11, 145.68, 146.59, 146.72, 147.41, 148.89, 154.00, 160.59, 170.20, 170.20, 170.20, 170.20.

4-(4-Methoxybenzyl)-3-(4-((2S,3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yl)amino) phenyl) quinazolin-4(3H)-one (11): m.p. 158-160 °C; yield 48 %. Anal. for C₂₈H₂₉N₃O₇ (m.w. 519): Calcd: C, 64.73; H, 5.58; N, 8.09; Found: C, 64.73; H, 5.60; N, 8.03. IR ν (cm⁻¹) 1662 (C=O), 3268, 3384 (OH bonded and non-bonded); MS: m/z (int. %) [M⁺] 519(25.2); ¹H-NMR (DMSO-d₆) δ 3.22 (s, 3H, OCH₃), 4.00 (s, 1H, sec NH), 4.37 (s, 2H, CH₂Bz), 3.29-3.87 (m, 6H, H-2, H-3, H-4, H-5, H-6a, H-6b), 3.58 (m, 4H, 2'-OH, 3'-OH, 4'-OH), 3.65 (s, 6'-OH), 5.01 (d, 1H, Hanom), 7.30-7.44 (m, 8H, Ar-H), 7.32- 8.19 (m, 4H, qui); ¹³C-NMR: 15.01, 61.89, 64.89, 71.21, 77.38, 74.11, 82.59, 91.62, 117.9, 117.9, 120.78, 121.11, 122.4, 122.4, 126.6, 126.70, 127.29, 127.79, 127.92, 130.78, 131.11, 132.4, 132.4, 133.39, 143.18, 146.90, 154.0, 160.59.

2-(3,3'-Dimethyl-[1,1'-biphenyl]amino-4-yl)-4-(4-methoxybenzyl)-4'-((1S,2S,3S,4R,5R)-2,3,4-trihydroxy-6-(hydroxymethyl)tetrahydro-2H-pyran-2-yl)- quinazolin-4(3H)-one (12): m.p. 188-190 °C; yield 48 %. Anal. for C₃₇H₃₇N₃O₆ (m.w. 619): Calcd: C, 71.72; H, 5.97; N, 6.78; Found: C, 71.69; H, 5.94; N, 6.73. IR ν (cm⁻¹) 1669(C=O), 3266, 3382(OH bonded and non-bonded); MS: m/z (int. %) [M⁺] 619(34.2); ¹H-NMR (DMSO-d₆) δ 2.34(s, 6H, 2CH₃), 3.32 (s, 3H, OCH₃), 4.0(s, 1H, sec NH), 4.47 (s, 2H, OCH₂Bz), 3.33-3.67 (m, 6H, H-2, H-3, H-4, H-5, H-6a, H-6b), 3.58 (m, 3H, 2'-OH, 3'-OH, 4'-OH), 3.65 (s, 1H, 6'-OH), 5.38 (d, 1H, H_{anom}), 6.90- 7.99 (m, 10H, Ar-H), 7.29 - 8.19 (m, 4 H, qui); ¹³C-NMR: 15.01, 17.61, 7.89, 61.92, 64.90, 71.18, 74.11, 77.38, 82.60, 91.90, 113.89, 120.78, 122.01, 125.11, 125.68, 126.59, 126.72, 126.72, 127.29, 130.00, 131.00, 131.62, 133.41, 134.39, 134.78, 137.20, 140.78, 142.01, 145.11, 145.68, 146.59, 146.72, 146.72,147.41, 148.89, 154.00, 160.59.

Synthesis of (E)-4-((4-Aminophenyl) sulfonyl)-N-(1-(4-methoxyphenyl) ethylidene) aniline (15): A mixture of dapsone **13** and *p*-methoxyacetophenone **14** (0.01 mol each) in 20 mL boiling absolute ethanol was heated under reflux for 4h. The resulting mixture was cooled and poured onto ice/water. The solid that separated out was filtered, washed, dried and crystallized from ethanol giving colorless crystals; m.p. 176-178 °C; yield 63 %; Anal. for C₂₁H₂₀N₂O₃S (m.w. 380): Calcd: C, 66.31; H, 5.26; N, 7.36; S, 8.42; Found: C, 66.30; H, 5.25; N, 7.31; S, 8.37. IR ν (cm⁻¹) 1160 (S=O), 1610 (C=N), 2993 (CH) and 3430 (NH); MS:


m/z (int. %) [M^+] 380 (24.4), 382 (5.5), 383(9.1); 1H -NMR (DMSO- d_6) δ 2.27 (s, 3H, CH_3), 3.94 (s, 3H, OCH_3), 6.27 (s, 2H, NH_2), 6.99 - 7.88 (m, 12H, 3 ArH).

4-(4-Methoxy)benzyl-N-(4-((1-(4-methoxy phenyl)ethylidene)amino)phenyl sulfonyl) phenyl) quinazolin-1-ylamine (17) : 4-Chloro-2-methoxybenzylquinazoline **16** and product **15** (0.01 mol each) in 20 mL absolute ethanol was heated under reflux for 6 h. The resulting mixture was cooled and poured onto ice / water. The solid that separated out was filtered, washed with water, air-dried and crystallized from ethanol to afford white crystals of **7**; m.p. 226-228 °C; yield 58 %; Anal. for $C_{37}H_{32}N_4O_4S$ (m.w. 628): Calcd: C, 70.70; H, 5.09; N, 8.91; S, 5.09; Found: C, 70.71; H, 5.07; N, 8.88; S, 5.03. IR ν (cm^{-1}) 1158 (S=O), 1619 (C=N), 2992 (CH) and 3330 (NH); MS: m/z (int. %) [M^+] 628(13.22), 552(100), 553(34.0), 554(6.7), 553(2.3), 555(1.6); 1H -NMR (DMSO- d_6) δ 2.27 (s, 3H, CH_3), 3.57, 3.94 (2s, 6H, $2OCH_3$), 4.00 (s, 1H, sec NH), 4.26 (s, 2H, $-CH_2$), 7.25-8.08 (m, 16H, Ar-H), 7.46-8.81 (m, 4H, qui); ^{13}C -NMR: 14.79, 18.41, 55.80, 60.52, 111.89, 114.42, 114.38, 114.52, 114.52, 123.31, 123.31, 124.19, 127.49, 127.60, 128.69, 128.69, 129.11, 129.11, 129.62, 129.62, 131.41, 132.02, 132.49, 133.43, 135.32, 136.32, 137.35, 138.73, 139.90, 140.43, 145.91, 150.91, 156.29, 162.88, 165.31, 171.42, 192.80.


N-4-(4-Methoxy)benzylphthalazin-1-yl)-N-(4-((1-(4-methoxyphenyl) ethylidene) amino) phenyl) sulfonyl) phenyl)furan-2-carboxamide (18): A mixture of Schiff base **17** (0.01 mol) and the 2-furoyl chloride (0.01 mol) was refluxed in 50 mL of dry pyridine for 3h. The excess solvent was distilled off and the reaction solution was cooled, then poured into crushed ice with frequent stirring leaving a crude product which was filtered off, washed with water, dried and crystallized from ethanol; m.p. 264-265 °C; yield 78%. Anal. for $C_{42}H_{34}N_4O_6S$ (m.w. 722): Calcd: C, 69.80; H, 4.70; N, 7.75; S, 4.43; Found: C, 69.78; H, 4.69; N, 7.74; S, 4.39. IR ν (cm^{-1}) 1157 (S=O), 1588 (C=N), 1734 (C=O), and 2988 (CH); MS: m/z (int.%) [M^+] 722(23.2), 646(18.2), 119(100); 1H -NMR(DMSO- d_6) δ 2.21 (s, 3H, CH_3), 3.35, 3.95 (2s, 6H, $2OCH_3$), 4.52 (s, 2H, CH_2), 7.28-8.33 (m, 16H, 3ArH), 7.24, 7.67, 8.41 (3dd, 3H, 3CH of furan), 7.55-8.83 (m, 4H, qui); ^{13}C -NMR: 14.80, 18.39, 55.82, 60.49, 111.68, 111.92, 114.40, 114.4, 115.3, 122.60, 122.60, 123.29, 123.29, 124.21, 127.48, 127.6, 128.52, 128.52, 128.68, 129.61, 129.61, 130.25, 131.54, 132.0, 132.49, 134.67, 136.34, 137.0, 139.92, 140.78, 143.78, 147.01, 147.71, 148.24, 150.89, 156.32, 157.68, 162.9, 165.31, 171.39, 172.34, 192.78.

RESULTS AND DISCUSSION

The reaction of benzoxazin-4-one with binucleophiles afforded quinazolinone derivatives that were highly antimicrobial effects[18, 27]. Quinazolinone could be synthesized according to our previously published work[28-42]. Quinazolinone inhibitors of inosine-5-monophosphate[43] that containing methylene, phenyl and methoxy groups. So, the authors can be decided the reaction of 3,2-benzoxazin-4-one with binucleophiles e.g. *p*-phenylenediamine, 4,4-benzidene, and 2,2-dimethyl-4,4-benzidene afforded phthalazinone derivatives **1-3** synthesis of some novel quinazolinone-containing Schiff bases, with the aim of obtaining more precise information about the course of reaction. Thus, 2-(4-aminophenyl)-4-methoxy benzylquinazolin-4(3H)-one(**1**), 2-(4'-amino-[1,1'-biphenyl]-4-yl)-4-methoxy-benzylquinazolin-4(3H)-one(**2**) and 2-(4'-amino-3,3'-dimethyl-[1,1'-biphenyl]-4-yl)-4-methoxy-N benzylquinazolin-4(3H)-one(**3**), each was reacted with *p*-methoxybenzaldehyde in boiling absolute ethanol giving products **4**, **5** and **6** respectively (Scheme 1). The IR spectra of **4**, **5** and **6** devoid any bands for the NH_2 group and the mass spectra showed molecular ion peaks at m/z 475, 551, and 579 for derivative **4**, **5** and **6** respectively.



On the other hand, when the amino quinazolinone derivative **4** and/or **5** was allowed to react with phenylisothiocyanate afforded azatedine derivative **7** and/or **8** respectively [36], (scheme 1). Moreover, The reaction of quinazolinone derivatives **1** and/or **2** with α -bromoglucose tetraacetate in 1,4-dioxane gave the per-*O*-acetylated quinazolinone **9**, and **10**, whose IR spectrum showed two strong peaks at 1668 and 1736 attributable for ν_{\max} of the two C=O groups of quinazolinone and peracetyl moieties, respectively, and the ¹H-NMR spectrum showed a singlet at δ 2.05-2.06 which is attributable for CH₃ of acetyl groups. The deacetylation of **9**, and **10** by sodium carbonate and ethanol gave the deacetylated derivative **11**, and **12**. The structure of **11** was assigned by its mass spectrum that showed a molecular ion peak at m/z 519 (M⁺). The authors can be reported also, the overall reaction in Scheme 2. Dapsone was refluxed with an equimolar amount of *p*-methoxyacetophenone in boiling ethanol giving the Schiff base **15** in good yield. The formation of product **15** was confirmed by its mass spectrum showing a molecular ion peak at m/z 380[M⁺] and its IR spectrum showing an absorption band at 1160 attributable for ν_{\max} of S=O in dapsone. The latter was reacted with 4-chloro-4-methoxybenzyl quinazoline **16** in boiling ethanol gave product **17** (Scheme 2). This product could be isolated by washing the crude reaction mixture with water, to remove any ammonium salts formed, then crystallized from ethanol. Purification of product **17** was rechecked by TLC. Formation of product **17** was confirmed by mass spectrum that showed a molecular ion peak at m/z 628(M⁺). In its pure state, product **7** was reacted with an equimolar amount of 2-furoyl chloride in dry pyridine affording derivative **18**, whose structure was inferred by the IR spectrum that showed a strong peak at 1734 attributable for ν_{\max} of the C=O attached to the furoyl moiety. Also, the IR devoid any band for *sec*-NH. The chlorine test for **18** showed no indication for the presence of chlorine.



APPLICATIONS

Biological investigation (Antimicrobial activity): The antimicrobial screening of all the synthesized compounds was done using the agar diffusion assay. This screening was performed against the Gram-positive bacteria, Gram-negative bacteria, staphylococcus aureus atcc 06538, Escherechia coli Atcc 10536, pathogenic fungi candida albicans Atcc 1023 and Aspergills Flavus. A moderate activity was observed with compounds which proved to possess marked activity against E. coli, S. aureus and C. albicans. The strong activity has observed nearly with the most synthesized compounds. The inhibitory concentration was determined for each of the active compounds along with Ampicillin, Streptomycin and Nystatin as positive control. No activity was detected for all the synthesized compound, toward Aspergillus flavus. Results are shown in the following table 1.

Table 1. Antimicrobial screening results of tested compounds at 1 mg 1 mL⁻¹

Compound No.	E. coli	S. aureus	A. Flavus	C. albicans
1	14	16	14	12
2	15	12	10	11
3	12	12	16	12
4	15	14	17	11
5	13	16	11	12
6	11	14	18	12
7	13	13	15	21
8	18	21	11	10
9	16	11	14	11
10	12	18	10	12
11	12	15	11	12
12	16	12	13	11
15	12	17	12	10
16	14	13	15	11
17	16	16	12	13
18	15	14	14	12
Ampicillin	0.0	22	0.0	0.0
Streptomycin	20	21	0.0	0.0
Nystatin	0.0	0.0	0.0	22

No activity (0.0), inhibition zone (< 7 mm), weak activity (7-10), moderate activity (11-15 mm), strong activity (> 15 mm), solvent CDCl₃ (6 mm).

REFERENCES

- [1] R. P. Jain, J. C. Vederas, *Bioorg. Med. Chem. Lett.*, **2004**, 14, 3655.
 [2] Y. Xu, Q. Guo, *Heterocycl*, **2004**, 63, 903.

- [3] G. Strappaghetti, C. Brodi, G. Giannaccini, L. Betti, *Bioorg. Med. Chem. Lett*, **2006**, 16, 2575.
- [4] G. W. Gribble, In *Comprehensive Heterocyclic Chemistry II*; A.R. Katrizky, C.W. Rees, E. V. Scriven, Eds.; Elsevier: Oxford, **1996**, 2, 207.
- [5] I. E. El-Shamy, A.M. Abdel-Mohsen, A. A. Alsheikh, M. G. Fouda, S. S. Al-Deyab, M. A. El-Hashash, J. Jancar, *Dyes and Pigments*, **2015**, 113, 357.
- [6] M. E. Wagdy, S. I. Hany, A. A. Hatem, N. F. Noha, M. Y. Mohieldin, *Euro. J. Med. Chem*, **2015**, 89, 549.
- [7] F. Alo-Asser, K.N. Zelenin, E. E. Lesiovskaya, I. P. Bezhan, B.A. Chakchir, *Pharm. Chem. J*, **2002**, 36, 598.
- [8] R. P. Jain, J. C. Vederas, *Bioorg. Med. Chem. Lett*, **2004**, 14, 3655.
- [9] R. W. Carling, K. W. Moore, L. I. Street, D. Wild, C. Isted, P. D. Leeson, S. Thomas, D. O-Conner, R. M. Mc-Kernan, K. Quirk, S. M. Cook, J. R. Atack, K. A. Waftord, S. A. Thompson, G. R. Dawson, P. Ferris, J. I. Castro, *J. Med. Chem*, **2004**, 47, 1807.
- [10] S. Grasso, G. De-Sarro, N. Micale, M. Zappala, G. Puia, M. Baraldi, C. J. Demicheli, *Med. Chem*, **2000**, 43, 2851.
- [11] A. Haikal, E. El-Ashry, J. Banoub, *Carbohydr. Res*, **2003**, 338, 2291.
- [12] J.S. Kim, H. Lee, M. Suh, H. P. Choo, S.K. Lee, H.J. Park, *Bioorg. Med. Chem*, **2004**, 12, 3683.
- [13] S. Demirayak, A. Karaburun, R. Beis, *Eur. J. Med. Chem*, **2003**, 39, 1089.
- [14] M. Johsen, K. Rehse, H. Petz, J. Stasch, E. Bischoff, *Arch. Pharmacol.*, **2003**, 336, 591.
- [15] E. Lenz, I. Wilson, B. Wright, E. Partidge, C. Roddgers, P.R. Haycock, *Pharm. Biomed. Anal*, **2002**, 28, 31.
- [16] E. Olmo, M. Armas, J. Lopez-perez, G. Ruiz, F. Vagas, A. Gimenez, *Bioorg. Med. Chem. Lett*, **2001**, 11, 2755.
- [17] D. Dogruer, E. Kupeli, E. Yesilada, M. Sahin, *Arch. Pharmacol*, **2004**, 337, 303.
- [18] Kh.M. Shakhidoyatov, in *Azotistye deterotsikly I alkaloidy [Nitrogen Heterocycles and Alkaloids]*, Eds V.G. Kartsev, G.A. Tolstikov, Iridium-Press, Moscow, 186 (in Russian), **2001**.
- [19] Z.Y. Zhang, C. H. Chu, X.P. Hui, *Ind. J. Chem.* **2002**, 41B, 2176.
- [20] H. Singh, M.K. Srivastava, B.K. Singh, S.K. Singh, G. Dubey, *Ind. J. Chem*, **2001**, 40B, 159.
- [21] K. Mogilaiah, S. Sakram, *Ind. J. Chem.* **2004**, 43B, 20 14.
- [22] M. Ajitha, K. Rajnarayana, K. Sangarapani, *Pharmazie*, **2002**, 57, 796.
- [23] S. Bhatk, M.S. Karthikeyan, B.S. Holla, N.S. Shetty, *Ind. J. Chem*, **2004**, 43B, 1765.
- [24] A.S. Aboraia, H.M.R. Rahman, N.M. Mahfouz, A.E.L. Gendy, *Bio-org. Med. Chem.* **2006**, 14, 1236.
- [25] S. Saxena, M. Verma, *Ind. J. Pharm. Sci.*, **1992**, 54, 1.
- [26] D. Mullican, M.W. Wilson, D.T. Cannor, C.R. Kostlan, D.J. Schirier, R. D. Dyer: *J. Med. Chem*, **1993**, 36, 1090.
- [27] P. Mishra, K. Joshi, *Ind. J. Physiol. Pharmacol*, **1992**, 36, 347.
- [28] A. M. F. Eissa, A. M. Elmetwally, M. A. Elhashash, A. M. Elgohary, *J. Korean Chem. Soc*, **2008**, 52, 3.
- [29] A. F. Fahmy, M., M. A. El-Hashash, M. M. Habishy, S. Nassar, *J. Revue Roumaine de Chimie*, **1978**, 23(11-12), 1567.
- [30] M. A. El-Hashash, K. M. Darwish, S. A. Rizk, F. A. El-Bassiouny; The Uses of 2-Ethoxy-4(H)-3,1-benzoxazin-4-one in the Synthesis of Some quinazolinone derivatives of antimicrobial Activity. *pharmaceuticals*, **2011**, 4, 1032-1051.
- [31] M. A. El-Hashash, K. M. Darwish, S. A. Rizk, F. A. El-Bassiouny; The reactivity of 2-Ethoxy-4-Chloroquinazoline and Its Use in Synthesis of Novel Quinazoline Derivatives ; *Organic .Chem. Intern.* **2011** ;doi;10.1155 /2011/295491.
- [32] Freddy H. Havaladar , Abhay R. Patil, Syntheses of 1, 2, 4 Triazole Derivatives and their Biological Activity, *E-Journal of Chemistry*, CODEN ECJHAO ISSN: 09734945;5,.2,.347354, **2008**.
- [33] Bing Chai, Xuhong Qian, Song Cao, Haidong Liu, Gonghua Song, Synthesis and insecticidal of 1,2,4-triazole derivatives, *ARKIVOC(II)* 141-145, **2003**.

- [34] S.I.El-Nagdy, M.A.El-Hashash, A.S.El-Badaway, Synthesis of New 4(3H)-Quinazolinone derivatives. *Egypt. J.Chem.* **2006**, 49(6),721-730.
- [35] M.S.Amine, M.A.El-Hashash, I.A.Attia; Synthesis and reaction of 2-ethoxy carbonyl-4(3H)-quinazolinone with nitrogen nucleophiles, *Indian J.Chem.* **1993**, 32B,577.
- [36] M. A. El-Hashash, S . A. Rizk, F. A. El-Bassiouny, K . M..Darwish; Syntheses of Novel Schiff bases and *N*-Nucleosides Bearing 2-Ethoxy quinazolin-4(3H)-one-3-yl or 2-Ethoxy quinazolin-4-yl Moieties, *Int. J. Chem. Sci. Tech.* **2011**, 1(4), 150-157.
- [37] M. A. El-Hashash, D. B. Guirguis, M. A. Kadhim, *J of American Science*, **2013**, 9, 12.
- [38] M. A. El-Hashash, S. A. Rizk, Nessim M.I, Utility of Benzimidazoles in Synthesis of New Bases of Nucleoside Moieties, and as Antioxidant in Lubricant Oils, *J Chem Eng Process Technol*, **2013**, 4,6.
- [39] M. A. El-Hashash, S. A. Rizk, A. Y. El-Kady, A. M. Khalifa; Design, Synthesis and Structural Characterisation of Some Phthalazin-1(2H)-one Derivatives as reactive dye, anticorrosive, and antimicrobial nucleosides; *J. Braz. Chem. Soc.*(Accepted), **2015**.
- [40] M. A. El-Hashash, S. A. Rizk, Sythesis of Some New Quinazolin-4-one Derivatives and Evaluation of Their Antimicrobial effects; *Egypt. J. Chem.* **2011**, 54, 4.
- [41] M. A. El-Hashash, S. A. Rizk, F. A. El-Bassiouny, K. M. Darwish; Uses of 2-Ethoxy-4(3H) quinazolinone in Synthesis of Quinazoline and Quinazolinone Derivatives of Antimicrobial Activity: The Solvent Effect, *Global J. Health Sci*, **2012**, 4(1).
- [42] M. A. El-Hashash, S. A. Rizk, Synthesis and Utility of 6-phthalimidoethyl-2(methyl /phenyl) benzoxazin-4-one in some heterocyclic synthesis; *Mid. E. J. Sci. Res*, **2012**, 11(4), 541.
- [43] C. R. Johnson, S. K. Gorla, M. Kavitha, M. Zhang, X. Liu, B. Striepen, J. R. Mead, G. D. Cuny, L. Hedstrom; Phthalazinone inhibitors of inosine-50-monophosphate dehydrogenase from *Cryptosporidium parvum*, *Bioorganic & Medicinal Chemistry Letters*, **2013**, 23, 1004.

AUTHORS' ADDRESSES

1. **M. A. EL-Hashash**
Department of Organic Chemistry,
Faculty of Science, University of Ain Shams,
Cairo, EGYPT
2. **M. A. Kadhim**
Department of Chemistry,
Faculty of Educational Science,
University of Anbar, IRAQ
Email- mak.organic@yahoo.com
3. **S. A. Rizk**
Department of Chemistry,
Facility of science,
University of Ain Shams, Cairo, EGYPT